

Math 333 Test 1
Solutions

1

1 a) $r^3 - 3r^2 + 4r - 2 = (r-1)(r^2 - 2r + 2) = 0$ has roots $r=1$ and $r = 1 \pm i$. Thus the general solution is

$$y = Ae^x + e^x(B\cos x + C\sin x)$$

b) $r^5 + 6r^4 + 12r^3 + 8r^2 = r^2(r^3 + 6r^2 + 12r + 8) = r^2(r+2)^3$, so $r=0$ with multiplicity 2 and $r=-2$ with multiplicity 3. Thus, the general solution is

$$y = Ax + B + (Cx^2 + Dx + E)e^{-2x}$$

2 a) $r^4 + 4r^2 = r^2(r^2 + 4)$ so $r=0$ occurs with multiplicity 2 and $r=2i$ occurs with multiplicity 1. $e^{2ix} \cos 2x$ corresponds to $\lambda = 2 \pm 2i$, which has multiplicity 0. So $y_p = e^{2ix}((Ax+B)\cos 2x + (Cx+D)\sin 2x)$

b) $r^5 - 2r^2 = r^2(r^3 - 2)$ so $r=0$ occurs with multiplicity 2. x^3 corresponds to $\lambda=0$, so

$$y_p = x^2(Ax^3 + Bx^2 + Cx + D)$$

c) $r^4 + 16 = 0$ gives $r^2 = \pm 4i \Rightarrow r = 2(\pm \frac{1}{\sqrt{2}} \pm \frac{i}{\sqrt{2}}) = \pm \sqrt{2} \pm i\sqrt{2}$. $e^{\sqrt{2}ix} \cos \sqrt{2}x$ corresponds to $\lambda = \sqrt{2} \pm i\sqrt{2}$, so this occurs with multiplicity 1. Thus $y_p = x e^{\sqrt{2}ix}((Ax+B)\cos \sqrt{2}x + (Cx+D)\sin \sqrt{2}x)$

3

$$\begin{aligned} y_1 &= x \\ y_1' &= 1 \\ y_1'' &= 0 \end{aligned}$$

$$\begin{aligned} y_2 &= x \sin x \\ y_2' &= \sin x + x \cos x \\ y_2'' &= 2 \cos x - x \sin x \end{aligned}$$

$$\begin{aligned} y_3 &= x \cos x \\ y_3' &= \cos x - x \sin x \\ y_3'' &= -\sin x - x \cos x \end{aligned}$$

$$y_1' x + y_2' x \sin x + y_3' x \cos x = 0$$

$$y_1' \cdot 1 + y_2' (\sin x + x \cos x) + y_3' (\cos x - x \sin x) = 0$$

$$y_1' \cdot 0 + y_2' (2 \cos x - x \sin x) + y_3' (-\sin x - x \cos x) = x$$

$$\begin{array}{ccc|c} x & x \sin x & x \cos x & 0 \\ 1 & \sin x + x \cos x & \cos x - x \sin x & 0 \\ 0 & 2 \cos x - x \sin x & -\sin x - x \cos x & x \end{array}$$

$$\rightarrow \begin{array}{ccc|c} 0 & -x^2 \cos x & x^2 \sin x & 0 \\ 1 & \sin x + x \cos x & \cos x - x \sin x & 0 \\ 0 & 2 \cos x - x \sin x & -\sin x - x \cos x & x \end{array}$$

$$\rightarrow \begin{array}{ccc|c} 1 & \sin x + x \cos x & \cos x - x \sin x & 0 \\ 0 & 1 & -\sin x / \cos x & 0 \\ 0 & 2 \cos x - x \sin x & -\sin x - x \cos x & x \end{array}$$

$$\rightarrow \begin{array}{ccc|c} 1 & 0 & \cos x + \sin^2 x / \cos x & 0 \\ 0 & 1 & -\sin x / \cos x & 0 \\ 0 & 0 & -x \cos x - x \sin^2 x / \cos x & x \end{array}$$

$$\rightarrow \begin{array}{ccc|c} 1 & 0 & 1/\cos x & 0 \\ 0 & 1 & -\sin x / \cos x & 0 \\ 0 & 0 & -x/\cos x & x \end{array} \rightarrow \begin{array}{ccc|c} 1 & 0 & 1/\cos x & 0 \\ 0 & 1 & -\sin x / \cos x & 0 \\ 0 & 0 & 1 & -\cos x \end{array}$$

$$\rightarrow \begin{array}{cccc} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -\sin x \\ 0 & 0 & 1 & -\cos x \end{array}$$

$$u_1' = 1, u_2' = -\sin x, u_3' = -\cos x \quad u_1 = x, u_2 = \cos x, u_3 = -\sin x$$

$$y_p = x^2 + x \sin x \cos x - x \sin x \cos x = x^2.$$

4/ $x^2 \cos 2x$ corresponds to $\lambda = 2i$ with multiplicity 3
 e^x corresponds to $\lambda = 1$ with multiplicity 0
 x^6 corresponds to $\lambda = 0$ with multiplicity 7

So the characteristic eq of smallest degree is

$$r^7(r-1)((r+2i)(r-2i))^3 = r^7(r-1)(r^2+4)^3$$

$$= r^7(r-1)(r^6 + 12r^4 + 48r^2 + 64)$$

$$= r^7(r^7 - r^6 + 12r^5 - 12r^4 + 48r^3 - 48r^2 + 64r - 64)$$

Solving ODE $y^{(14)} - y^{(13)} + 12y^{(12)} - 12y^{(11)} + 48y^{(10)} - 48y^{(9)} + 64y^{(8)} - 64y^{(7)} = 0$

3 (Using determinants)

$$W = \begin{vmatrix} x & x \sin x & x \cos x \\ 1 & \sin x + x \cos x & \cos x - x \sin x \\ 0 & 2 \cos x - x \sin x & -2 \sin x - x \cos x \end{vmatrix}$$

$$= x \left(-2 \sin^2 x - 3x \cos x \sin x - x^2 \cos x - (2 \cos^2 x - 2x \sin x \cos x - x \sin x \cos x + x^2 \sin^2 x) \right)$$

$$= \left(-2x \sin^2 x - x^2 \sin x \cos x - 2x \cos^2 x + x^2 \sin x \cos x \right)$$

$$= -2x \sin^2 x - 3x^2 \sin x \cos x - x^3 \cos x - 2x \cos^2 x + 2x^2 \sin x \cos x + x^2 \sin x \cos x - x^3 \sin^2 x + 2x \sin^2 x + x^2 \sin x \cos x + 2x \cos^2 x - x^2 \sin x \cos x$$

$$= -x^3$$

$$W_1 = \begin{vmatrix} 0 & x \sin x & x \cos x \\ 0 & \sin x + x \cos x & \cos x - x \sin x \\ 1 & 2 \cos x - x \sin x & -2 \sin x - x \cos x \end{vmatrix}$$

$$= x \sin x \cos x - x^2 \sin^2 x - x \sin x \cos x - x^2 \cos^2 x$$

$$= -x^2$$

$$W_2 = \begin{vmatrix} x & 0 & x \cos x \\ 1 & 0 & \cos x - x \sin x \\ 0 & 1 & -2 \sin x - x \cos x \end{vmatrix}$$

$$= - (x \cos x - x^2 \sin x - x \cos x)$$

$$= x^2 \sin x$$

$$W_3 = \begin{vmatrix} x & x \sin x & 0 \\ 1 & \sin x + x \cos x & 0 \\ 0 & 2 \cos x - x \sin x & 1 \end{vmatrix}$$

$$= x \sin x + x^2 \cos x - x \sin x = x^2 \cos x$$

$$y_p = y_1 \int \frac{W_1}{W} r(x) + y_2 \int \frac{W_2}{W} r(x) + y_3 \int \frac{W_3}{W} r(x)$$

$$= x \int \frac{-x^2}{-x^3} \cdot x dx + x \sin x \int \frac{x^2 \sin x}{-x^3} \cdot x dx + x \cos x \int \frac{x^2 \cos x}{-x^3} \cdot x dx$$

$$= x \int 1 dx + x \sin x \int -\sin x dx + x \cos x \int -\cos x dx$$

$$= x^2 + x \sin x \cos x - x \cos x \sin x =$$

$$= x^2$$